

**List of commodities reported as potential hosts of Fall Armyworm**

Host	Host
<i>Allium sp.</i>	<i>Panicum</i> (millets)
<i>Allium cepa</i> (onion)	<i>Pennisetum glaucum</i> (pearl millet)
<i>Allium sativum</i> (garlic)	<i>Phaseolus vulgaris</i> (common bean)
<i>Arachis hypogaea</i> (groundnut)	<i>Pisum sativum</i> (pea)
<i>Asparagus officinalis</i> (asparagus)	<i>Prunus persica</i> (peach)
<i>Beta sp.</i>	<i>Pyrus communis</i> (European pear)
<i>Brassica sp.</i>	<i>Raphanus sativus</i> (radish)
<i>Capsicum sp.</i>	<i>Rosa</i> (roses)
<i>Carya illinoiensis</i> (pecan)	<i>Sesamum indicum</i> (sesame)
<i>Chenopodium quinoa</i> (quinoa)	<i>Setaria italica</i> (foxtail millet)
<i>Cicer arietinum</i> (chickpea)	<i>Setaria parviflora</i> (knot root foxtail)
<i>Citrullus lanatus</i> (watermelon)	<i>Solanum lycopersicum</i> (tomato)
<i>Citrus sp.</i>	<i>Solanum tuberosum</i> (potato)
<i>Coffea arabica</i> (arabica coffee)	<i>Sorghum</i>
<i>Cucumis melo</i> (melon)	<i>Spinacia deracea</i> (spinach)
<i>Cucumis sativus</i> (cucumber)	<i>Trifolium</i> (clovers)
<i>Cucurbitaceae</i>	<i>Trifolium incarnatum</i> (Crimson clover)
<i>Fagopyrum esculentum</i> (buckwheat)	<i>Triticum</i> (wheat)
<i>Ficus</i>	<i>Triticum aestivum</i> (wheat)
<i>Fragaria sp.</i>	<i>Vaccinium</i> (blueberries)
<i>Glycine max</i> (soyabean)	<i>Vaccinium corymbosum</i> (blueberry)
<i>Gossypium</i> (cotton)	<i>Vicia faba</i> (faba bean)
<i>Helianthus annuus</i> (sunflower)	<i>Vigna unguiculata</i> (cowpea)
<i>Hordeum vulgare</i> (barely)	<i>Vitis</i> (grape)
<i>Ipomoea batatas</i> (sweet potato)	<i>Vitis vinifera</i> (grapevine)
<i>Lactuca sativa</i> (lettuce)	<i>Zea mays</i> (maize)
<i>Linum usitatissimum</i> (flax)	<i>Zea mays subsp. mays</i> (sweetcorn)
<i>Malus domestica</i> (apple)	Cut flowers and foliage